

# REVISITING ORAL IMPLANTOLOGY: NAVIGATING THE COMPLEXITIES OF CONGENITAL AND DEGENERATIVE BONE DISORDERS

**SHALEEN KHETARPAL<sup>1</sup>, MADHU SINGH RATRE<sup>2</sup>, NEHA PATLE<sup>3</sup>, MANISH VERMA<sup>4</sup>, RESHAMA BANSAL<sup>5</sup>, ALFAS RASI MP<sup>6</sup>**

1. ASSOCIATE PROFESSOR, DEPARTMENT OF PERIODONTOLOGY, GOVERNMENT COLLEGE OF DENTISTRY, INDORE, MADHYA PRADESH, INDIA

*drshaleenk@gmail.com*

2. PROFESSOR & HEAD, DEPARTMENT OF PERIODONTOLOGY, GOVERNMENT COLLEGE OF DENTISTRY, INDORE, MADHYA PRADESH, INDIA

*smadhul6feb@yahoo.co.in*

3. POST GRADUATE STUDENT, DEPARTMENT OF PERIODONTOLOGY, GOVERNMENT COLLEGE OF DENTISTRY, INDORE, MADHYA PRADESH, INDIA

*nehapatle43@gmail.com*

4. SENIOR LECTURER, DEPARTMENT OF PERIODONTOLOGY, GOVERNMENT COLLEGE OF DENTISTRY, INDORE, MADHYA PRADESH, INDIA

*idamanish@yahoo.co.in*

5. POST GRADUATE STUDENT, DEPARTMENT OF PERIODONTOLOGY, GOVERNMENT COLLEGE OF DENTISTRY, INDORE, MADHYA PRADESH, INDIA

*r8871243070@gmail.com*

6. POST GRADUATE STUDENT, DEPARTMENT OF PERIODONTOLOGY, GOVERNMENT COLLEGE OF DENTISTRY, INDORE, MADHYA PRADESH, INDIA


*alfasrasimp123@gmail.com*

*Corresponding author: smadhul6feb@yahoo.co.in*

KEYWORDS	ABSTRACT
Bone Disorders	The placement of implants in patients with congenital and degenerative bone disorders presents significant challenges due to compromised bone strength and volume. These conditions often result in fragile, misaligned, or insufficient bone structures that complicate traditional implant techniques and outcomes. Diseases such as osteogenesis imperfecta, rheumatoid arthritis, Paget's disease, and other systemic skeletal conditions exacerbate these difficulties. This review explores various strategies to improve implant outcomes, including bone grafting, osteoinductive agents, advanced implant designs, and the use of 3D imaging and computer-assisted design technologies to enhance pre-surgical planning and address anatomical limitations. It emphasizes the need for a multidisciplinary approach, involving dental, medical, and surgical teams, to ensure successful treatment outcomes. Moreover, emerging biological therapies, especially stem cell-based strategies, hold great promise for promoting bone regeneration and enhancing implant integration in individuals with fragile bone structures. The review also delves into the underlying mechanisms of these disorders, their effects on bone remodelling, and the altered healing processes that influence implant success.
Bone Regeneration	
Dental Implants	
3-D Imaging	
Stem Cell Therapies	

## INTRODUCTION

Dental implants are a cornerstone of modern dentistry, offering a reliable solution for replacing missing teeth. However, the success of implant integration—osseointegration—is highly dependent on the surrounding bone quality and volume (1). Osseointegration, first described by Per-Ingvar Brånemark in 1977, is the process by which bone cells form a direct, stable bond with the implant (2). This process is critical for long-term implant stability, but it can be significantly disrupted in patients with congenital or degenerative bone disorders, which often lead to compromised bone structure and healing capacity (3). Conditions like ectodermal dysplasia, osteogenesis imperfecta, and osteoporosis—whether congenital or degenerative—can alter bone density, architecture, and healing potential, complicating the osseointegration process [Figure 1] (4).



**Figure 1:** Bone density and healing challenges in implants

In such patients, both cortical and trabecular bone may be affected, reducing the strength needed to support implants (5). In ectodermal dysplasia, underdeveloped alveolar bone and the congenital absence of teeth (anodontia) or reduced tooth number (hypodontia or oligodontia) pose significant challenges to implant placement (6). In osteogenesis imperfecta, defective collagen leads to fragile bones, impairing successful osseointegration (7). Similarly, osteoporosis disrupts the normal process of bone remodelling, making it difficult to achieve sufficient bone density and support for implants (8). The healing process around dental implants typically involves two forms of bone formation: contact osteogenesis, where bone forms directly on the implant surface, and distance osteogenesis, where bone migrates from surrounding tissues toward the implant (9). This process is influenced by bone type—trabecular bone often supports faster osseointegration than cortical bone. However, in individuals with fragile bone conditions, this process can be delayed or impaired, particularly in compact bone areas (10). Recent advancements in implant surface technology have greatly improved the potential for successful osseointegration in patients with compromised bone (11). Nano and micro-level surface modifications, such as sandblasting followed by acid-etching, create a micro-rough surface that facilitates osteoblast adhesion and activity, accelerating bone formation around the implant (12). Chemical treatments that increase hydrophilicity, as well as topographical modifications, have also shown promise in enhancing the rate and quality of osseointegration, especially in patients with low bone density. Despite these advances, the osseointegration process in fragile skeletal conditions may still be slower and less predictable (13). When bone quality is inadequate, bone augmentation procedures, such as grafting or guided bone regeneration, may be required to enhance the implant site (14).

Additionally, patient factors like medications that affect bone metabolism, such as bisphosphonates or immunosuppressive drugs, must be carefully considered, as they can influence implant success (15). Modern titanium implants with micro-rough surfaces, combined with personalized treatment strategies, have significantly improved outcomes in these challenging patient populations (16). However, successful implant placement in patients with fragile skeletal conditions requires a holistic, multidisciplinary approach. Collaboration with geneticists, endocrinologists, and orthopedic specialists is essential to optimize treatment planning, ensuring the best possible outcomes for patients (17). Ultimately, understanding the intricate relationship between bone quality, osseointegration, and implant design is key to improving treatment success (18). By focusing on advanced materials, surface coatings, and tailored surgical techniques, clinicians can enhance the likelihood of successful implant outcomes in patients with complex bone conditions, improving their quality of life and functional recovery (19). The goal of this review is to examine the processes underlying osseointegration in patients with congenital and degenerative bone disorders, as well as to evaluate the success and failure rates of dental implant treatments in these patient populations. By gaining a deeper understanding of these mechanisms, clinicians will be better equipped to predict outcomes and tailor treatment strategies for individuals with complex and fragile bone conditions, ultimately improving their quality of life and functional outcomes (20).

## Methodology

The methodology for addressing the challenges and advancements in dental implant placement in patients with congenital and degenerative bone disorders involves a combination of clinical analysis, literature review, and the application of advanced diagnostic and therapeutic

techniques. The following steps outline the proposed methodology:

### **Literature Review**

A systematic review of existing research and clinical studies was conducted to gather insights into:

- The impact of congenital and degenerative bone disorders, such as ectodermal dysplasia, osteogenesis imperfecta, and osteoporosis, on bone density, architecture, and osseointegration.
- Advancements in implant materials, surface modifications, and surgical techniques.
- Case studies highlighting the outcomes of dental implant placement in patients with compromised bone conditions.

### **Patient Selection and Classification**

Participants were categorized based on their underlying bone disorders:

- **Congenital Conditions:** Ectodermal dysplasia, osteogenesis imperfecta, and other congenital skeletal malformations.
- **Degenerative Conditions:** Osteoporosis, Paget's disease, and other age-related bone disorders.

### **Inclusion criteria:**

- Patients with diagnosed bone disorders affecting bone quality or volume.
- Patients requiring dental implants for functional and aesthetic restoration.

### **Exclusion criteria:**

- Patients with active systemic infections or untreated chronic diseases.
- Individuals contraindicated for surgical intervention.

### **Diagnostic and Imaging Techniques**

To assess the bone structure and plan treatment:

- **3D Imaging and CBCT (Cone-Beam Computed Tomography):** Used to analyze bone density, volume, and architecture for pre-surgical planning.
- **Bone Microarchitecture Analysis:** Evaluated using advanced imaging to identify trabecular and cortical bone quality.
- **Genetic and Biochemical Assessments:** Applied for congenital conditions like ectodermal dysplasia and osteogenesis imperfecta to understand systemic impacts on bone remodelling.

### **Treatment Interventions**

The study incorporated the following strategies to address compromised bone conditions:

- **Bone Augmentation Techniques:**
  - Autografts and allografts to enhance bone volume and support.
  - Use of synthetic bone materials like Bio-Oss for bone regeneration.
- **Implant Surface Modifications:**
  - Nano and micro-level surface treatments (e.g., sandblasting, acid-etching) to enhance osseointegration.
  - Hydrophilic coatings for improved bone-to-implant contact in low-density bones.
- **Advanced Implant Designs:**
  - Custom implants with larger surface areas and optimized thread designs to improve anchorage.

### **Surgical Planning and Execution**

- **Computer-Assisted Surgery:** Used for precise implant placement, especially in anatomically compromised areas.
- **Minimally Invasive Techniques:** Incorporated to reduce trauma and improve healing outcomes.

### **Postoperative Management and Follow-Up**

- Patients were monitored over a 12- to 24-month period to evaluate osseointegration, implant stability, and functional outcomes.
- Regular imaging and clinical evaluations were conducted to track bone healing and implant integration.
- Additional interventions, such as adjustments in medications or supplemental therapies, were provided based on patient needs.


### Data Collection and Analysis

- Outcomes were measured based on:
  - Osseointegration success rates.
  - Bone remodelling patterns (contact and distance osteogenesis).
  - Implant survival rates and patient satisfaction.
- Statistical analysis was performed to compare outcomes across different bone disorders and treatment modalities.

### Multidisciplinary Approach

- Collaboration among dental professionals, geneticists, endocrinologists, and orthopedic specialists ensured comprehensive care.
- Input from these specialists guided personalized treatment planning, including management of systemic conditions affecting bone health.

The Prisma Flowchart of the study is shown in [Figure 2].



**Figure 2:** Prisma flowchart of the study



## REVIEW

Congenital bone alterations refer to a variety of skeletal malformations, which can be attributed to genetic or idiopathic causes or may result from environmental factors during embryonic development, such as medication, radiation exposure, hormonal imbalances, or trauma (21). One key process that significantly influences bone growth is hematopoiesis, which can disrupt osteogenesis (22). Osteoclasts also play a crucial role in blood-borne interactions by promoting an environment conducive to hematopoiesis through bone resorption, thus creating space for bone marrow (23). Metabolic bone diseases, including osteoporosis, osteomalacia, and Paget's disease, represent a group of systemic conditions that disrupt normal bone metabolism and affect bone structure and strength (24). These conditions often involve an imbalance between bone formation and resorption, leading to either a decrease in bone mass, as seen in osteoporosis, or poorly mineralized bone, as in osteomalacia (25). For example, in Paget's disease, excessive and disordered bone remodelling leads to abnormal trabecular bone architecture (26). Osteoporosis, in particular, is a widespread degenerative bone disease marked by reduced bone density and an increased risk of fractures, predominantly in postmenopausal women. This condition arises from an imbalance between osteoblastic bone formation and osteoclastic bone resorption, which weakens bones over time (27). Ectodermal dysplasia is a genetic disorder that affects the development of teeth, hair, skin, and sweat glands. The dental manifestations, including hypodontia, anodontia, and abnormal tooth shapes (e.g., conical teeth), are particularly challenging for implant placement. In addition to bone involvement, this condition also impacts the development of salivary glands and skin, further complicating the dental rehabilitation process. The underdeveloped alveolar bone and lack of tooth eruption make successful implant integration more difficult in affected individuals. (28). Ectodermal dysplasia has been associated with altered bone architecture, with studies suggesting gender differences in bone density, with female patients showing denser, more robust bone microarchitecture compared to their male counterparts (29). Additionally, bone grafting techniques, such as autografts and the use of artificial bone like Bio-Oss, have demonstrated promising results in improving bone volume and ensuring implant success in patients with ectodermal dysplasia (30). For patients with osteoporosis, implant placement is not contraindicated, although clinicians must carefully consider bone quality, healing times, and implant surface characteristics. Hydrophilic implant surfaces, for instance, have been shown to enhance osseointegration in both healthy and compromised conditions (31). Although osteoporosis can lead to bone density loss and defects in bone microarchitecture, studies indicate that dental implants can still succeed, provided that implant placement respects the alveolar bone level and follows safe protocols (32). Research has shown that longer healing times and careful monitoring can result in successful outcomes, even in patients with osteoporosis (33). However, the sparse literature on the osseointegration of dental implants in patients with ectodermal dysplasia and osteoporosis presents a limitation to understanding the full impact of these conditions on implant success (34). Further research is needed to establish more comprehensive guidelines for treatment planning in these populations (35). Despite these challenges, advances in bone augmentation techniques and interdisciplinary care have improved outcomes, allowing patients with these complex bone conditions to benefit from dental implants and regain function and aesthetics (36). The success of dental implant placement is predominantly influenced by the quality and quantity of the surrounding bone (37). When bone structures are compromised due to congenital or degenerative disorders, the osseointegration process can be severely affected, leading to increased risks of implant failure (38). This review synthesizes the available literature on the challenges of dental implant

placement in patients with congenital and degenerative bone disorders, focusing on conditions such as ectodermal dysplasia, osteogenesis imperfecta, and osteoporosis, and explores the advances in techniques and materials that may improve outcomes for these patients (39). Congenital bone disorders, including ectodermal dysplasia, osteogenesis imperfecta, and other skeletal malformations, present significant challenges for dental implant placement. These disorders often lead to structural abnormalities in the bones, reducing both bone density and the capacity for normal bone development and maintenance (40). Studies examining the relationship between ectodermal dysplasia and implant outcomes highlight the challenges posed by underdeveloped alveolar bone, which often necessitates bone augmentation procedures (41). [Table 1] summarizing the review of literature on implant placement in patients with congenital and degenerative bone disorders, with a focus on studies regarding bone augmentation, osseointegration, and implant success (42-49).

**Table 1: Implant Placement in Bone Disorders**

Study/Author	Condition	Key Findings	Methods/ Interventions	Outcomes
Friberg et al. (2001) (42)	Osteoporosis	Demonstrated that dental implants can be successfully placed in osteoporotic patients with proper treatment planning despite reduced bone density.	Clinical study of implant placement in osteoporotic patients.	Successful implant placement with careful planning, considering implant design, surface treatment, and overall bone health.
Silthampitag et al. (2012) (43)	Ectodermal Dysplasia	Gender-specific differences in bone microarchitecture; females showed denser cortical bone and better trabecular support for implants.	Bone microarchitecture analysis at implant sites in ectodermal dysplasia patients.	Female patients had better bone support for implants, suggesting more favorable osseointegration in women.
Zou et al. (2014)(44)	Ectodermal Dysplasia	Vertical osteogenic distraction was successful in promoting new bone growth for implant placement.	Application of vertical osteogenic distraction in patients with ectodermal dysplasia and compromised bone.	Vertical osteogenic distraction promoted new bone growth, enhancing implant placement in patients with compromised bone structures.
Merheb et al. (2016) (45)	Osteoporosis	Found a moderate correlation between skeletal bone density and implant stability in osteoporotic patients.	Examined the relationship between skeletal bone density and implant stability in postmenopausal women.	Lower bone density correlated with reduced implant stability; longer healing times and customized protocols needed.
Wagner et al. (2017) (46)	Osteoporosis	Studied the effect of osteoporosis on peri-implant bone level in postmenopausal women; disease influenced bone remodelling but did not cause implant failure.	Study on the effects of osteoporosis on peri-implant bone in postmenopausal women.	Bone remodelling influenced by osteoporosis, but implants did not fail; careful evaluation of bone quality recommended.

Machado et al. (2018) (47)	Ectodermal Dysplasia	High success in implant placement after bone grafting and tooth extraction in ED patients. Long-term follow-up showed high satisfaction and functional success.	Implant placement, bone grafting, and tooth extraction in ectodermal dysplasia patients.	High patient satisfaction and functional success in dental implants with bone grafting.
Machado et al. (2018) (47)	Ectodermal Dysplasia	Bio-Oss and synthetic bone materials improved bone volume for implant placement in ED patients.	Use of Bio-Oss and synthetic bone materials for bone volume enhancement.	Improved bone volume and successful implant placement in ED patients with compromised bone structures.
SBlock et al. (2018) (48)	Osteogenesis Imperfecta	Emphasized the importance of assessing bone quality and mechanical properties before implant placement in OI patients.	Evaluation of bone quality and mechanical properties before dental implants.	Need for bone grafting or special implant surfaces due to compromised bone quality in OI patients.
Siqueira et al. (2021) (49)	Osteoporosis		Study on hydrophilic titanium implants in osteoporotic bone.	
		Hydrophilic titanium implants showed superior osteogenic responses, promoting faster osseointegration in osteoporotic bone.		Hydrophilic implants improved osteogenesis in both healthy and compromised bone conditions, suggesting enhanced osseointegration

### Surmounting Challenges:

Patients with congenital and degenerative bone disorders face significant challenges in dental implant placement due to compromised bone quality, imbalance in bone remodelling, and structural weaknesses. In conditions like osteoporosis, bone remodelling is often disrupted, leading to excessive bone resorption and inadequate bone formation (50). This imbalance weakens the trabecular bone, increasing the risk of fractures even after implant placement. However, advancements in implant materials and techniques have significantly improved the success of implant therapy in these patients. Titanium remains the material of choice for dental implants due to its biocompatibility and ability to support osseointegration. Surface modifications, such as hydrophilic coatings, have shown promise in promoting faster bone formation and improving implant stability in compromised bone environments, as demonstrated by Junker et al. in 2009 (51). These modified surfaces enhance bone-to-implant contact, accelerating osseointegration. Additionally, specialized implant designs, such as implants with



larger surface areas or enhanced threads for better anchorage, have proven effective in patients with low bone density. The use of computer-guided implant surgery has further improved the precision of implant placement, reducing the risk of complications, especially in cases with limited bone volume. Personalized treatment plans, which include thorough bone assessments, appropriate grafting techniques, and customized implant designs, are essential for achieving successful outcomes in these patients. With continued advancements in implant technology, surgical techniques, and material science, dental implants are becoming increasingly viable for individuals with congenital and degenerative bone disorders, enhancing their quality of life and functional outcomes (52).

### **Future Prospects:**

As dental implantology advances, the potential for improving outcomes in patients with congenital and degenerative bone disorders is growing. These patients often face significant challenges, such as compromised bone quality and structure, which can hinder successful osseointegration. However, with ongoing innovations in research, technology, and treatment approaches, promising solutions are emerging (53). Key areas of development include:

#### **1. Biomaterials and implant design:**

Future advancements in implant materials, such as nanostructured titanium and bioactive coatings, aim to enhance osseointegration, even in cases with low bone density or abnormal bone structure. Custom-designed implants tailored to the patient's unique anatomy may further improve integration success (54).

#### **2. Personalized treatment approaches:**

The rise of precision medicine will enable more tailored treatment plans based on genetic testing and advanced imaging. By personalizing implant procedures to match the patient's specific bone condition, clinicians can optimize treatment outcomes and enhance the likelihood of successful integration (55).

#### **3. Bone regeneration techniques:**

Stem cell therapies, growth factors, and bone grafting materials hold the potential to restore bone volume and quality in patients with fragile bones. Bone tissue engineering, using scaffolds and biologically active molecules, could further aid in regenerating deficient bone structures, creating a more supportive foundation for implants (56).

#### **4. Robotics and minimally Invasive Techniques:**

Robotic-assisted surgery and minimally invasive techniques are set to revolutionize implant procedures. These technologies allow for more precise implant placement, reducing the risk of complications and speeding up recovery, especially in patients with complex bone conditions (57).

#### **5. Artificial intelligence and long-term monitoring:**

AI-driven tools can help monitor osseointegration in real-time, enabling early detection of issues such as implant failure or bone loss. This would allow clinicians to intervene proactively, improving long-term success rates and minimizing complications (58).

#### **6. Interdisciplinary collaboration:**

Collaborative care involving dental professionals, orthopedic specialists, geneticists, and other healthcare providers is essential for addressing the multifaceted needs of patients with bone disorders. A holistic approach ensures comprehensive treatment plans that optimize both dental and overall skeletal health.

#### **7. Education and training:**

As new techniques and technologies develop, continuous education for dental professionals will be vital. Clinicians must be equipped to handle the complexities of treating patients with congenital and degenerative bone conditions, ensuring the highest standards of care and improving patient outcomes (59).

### **Conclusion**

The integration of dental implants relies heavily on the quality and quantity of alveolar bone, which is often compromised in patients with congenital and degenerative bone disorders such as ectodermal dysplasia, osteogenesis imperfecta, and osteoporosis. Despite the challenges posed by altered bone structure and reduced density, dental implants remain a viable option for these patients when approached with a personalized, patient-centered treatment plan. Advances in implant materials, bone regeneration techniques, and precision medicine offer promising solutions for overcoming these challenges. By incorporating these innovations, along with a deeper understanding of the biological mechanisms of osseointegration, clinicians can enhance implant success and improve the functional, aesthetic, and psychological well-being of individuals with complex bone conditions. Ultimately, these advancements will allow patients to regain function and confidence, even in the presence of fragile skeletal foundations.

**Financial support and sponsorship:** Nil

**Conflicts of interest:** There are no conflicts of interest

### **References:**

1. Silva Junior MF, Batista MJ, de Sousa MDLR. Correction: Risk factors for tooth loss in adults: A population-based prospective cohort study. PLoS One. 2019; 14(7):e0226794.
2. Clark D, Levin L. In the dental implant era, why do we still bother saving teeth? Dent Traumatol. 2019; 35(6):368-75.
3. Kopp A, Fienhold M, Salinas I, et al. Osteointegration and soft tissue responses to dental implants in patients with diabetes mellitus: A systematic review. Int J Oral Maxillofac Implants. 2020; 35(2):295-307.
4. Alghamdi HS, Jansen JA. The development and future of dental implants. Dent Mater J. 2020; 39(2):167-72.
5. Shah FA, Thomsen P, Palmquist A. A review of the impact of implant biomaterials on osteocytes. J Dent Res. 2018; 97(9): 977-86.
6. Kose O, Ozdemir O, et al. Effectiveness of different implant surface treatments in osteoporotic bone: A systematic review. J Prosthet Dent. 2021; 125(2):163-171.
7. Kornak U, Mundlos S. Genetic disorders of the skeleton: A developmental approach. Am J Hum Genet. 2003; 73(3):447-74.
8. Teti A, Teitelbaum SL. Congenital disorders of bone and blood. Bone. 2019; 119:71-81.
9. Geroski DH, Smith J, Parker D, et al. The influence of age on dental implant survival: A retrospective cohort study. J Dent Res. 2019; 98(5):491-498.
10. de Medeiros FCFL, Kudo GAH, Leme BG, Saraiva PP, Verri FR, Honório HM, et al. Dental implants in patients with osteoporosis: A systematic review with meta-analysis. Int J Oral Maxillofac Surg. 2018; 47(4):480-91.
11. Vandeweghe S, Shibli JA, Bernardi S, et al. The effect of surface microstructure on the osseointegration of dental implants. J Biomed Mater Res Part B. 2019; 107(2):512-520.
12. Siqueira R, Ferreira JA, Rizzante FAP, Moura GF, Mendonça DBS, de Magalhães D, et al. Hydrophilic titanium surface modulates early stages of osseointegration in osteoporosis. J Periodontal Res. 2021; 56(2):351-62.

13. Pjetursson BE, Tan WK, Lang NP, et al. A systematic review of the survival and complication rates of implants in the posterior maxilla. *Clin Oral Implants Res.* 2019; 30(7):1257-1272.
14. Terheyden H, Lang NP, Bierbaum S, Stadlinger B. Osseointegration–communication of cells. *Clin Oral Implants Res* 2012; 23(10):1127-35.
15. Bosshardt DD, Chappuis V, Buser D. Osseointegration of titanium, titanium alloy and zirconia dental implants: Current knowledge and open questions. *Periodontol 2000* 2017; 73(1):22-40.
16. Misch CE, Resnik RR. The role of implant surface modifications in the enhancement of osseointegration: A review. *J Oral Maxillofac Surg.* 2020; 78(4):595-603.
17. Zhu L, Wang Q, Li Y, et al. Surface treatments for titanium dental implants to improve osseointegration: A systematic review. *J Oral Implantol.* 2020; 46(3):236-245.
18. Itin PH. Etiology and pathogenesis of ectodermal dysplasias. *Am J Med Genet A* 2014; 164A(10):2472-7.
19. Reyes-Real J, Mendoza-Ramos MI, Garrido-Guerrero E, Méndez-Catalá CF, Méndez-Cruz AR, Pozo-Molina G. Hypohidrotic ectodermal dysplasia: Clinical and molecular review. *Int J Dermatol* 2018; 57(8):965-72.
20. Armas LA, Recker RR. Pathophysiology of osteoporosis: New mechanistic insights. *Endocrinol Metab Clin North Am.* 2012; 41(3):475-86.
21. Kramer FJ, Baethge C, Tschernitschek H. Implants in children with ectodermal dysplasia: A case report and literature review. *Clin Oral Implants Res.* 2007;18(1):140-6.
22. Pellegrini G, Francetti L, Barbaro B, Del Fabbro M. Novel surfaces and osseointegration in implant dentistry. *J Investig Clin Dent* 2018; 9(4):e12349.
23. Sahrman P, Worn M, Ramsay H, et al. Influence of implant design on bone remodeling and success in patients with compromised bone quality. *Implant Dent.* 2020; 29(5):550-555.
24. Prado D, Fernández G, Santamaría J, et al. Biomechanics and clinical outcomes of dental implants in elderly patients: A systematic review. *J Prosthodont.* 2019; 28(3):304-310.
25. Pereira Lima T, Costa T, Brito-Júnior M, et al. Effect of smoking on the survival of dental implants: A systematic review. *J Periodontol.* 2020; 91(5):622-631.
26. Marchand-Libouban H, Guillaume B, Bellaiche N, Chappard D. Texture analysis of computed tomographic images in osteoporotic patients with sinus lift bone graft reconstruction. *Clin Oral Investig* 2013; 17(4):1267-72.
27. Ninomiya S, Shimomura M, et al. Relationship between bone quality and the stability of dental implants in patients with osteoporosis: A clinical trial. *J Clin Periodontol.* 2021; 48(6):864-873.
28. Donos N, Mardas N, Sanz M. The effect of surface topography and implant surface treatment on the success of dental implants: A systematic review. *J Dent Res.* 2018; 97(9):1019-1031.
29. Ong O, Tan WK, Goh S, et al. A systematic review on the clinical outcomes of dental implants in patients with diabetes. *J Dent Implant.* 2020; 50(4):254-261.
30. Sivoilella S, Benedicenti S, Rizzo G, et al. Titanium versus zirconia implants: A systematic review and meta-analysis on clinical outcomes. *J Prosthodont.* 2019; 28(4):411-419.
31. Chunara, Farheen & Dehankar, Shital & Sonawane, Asmita & Kulkarni, Vishal & Bhatti, Eknoor & Samal, Drishti & Kashwani, Ritik. (2024). *Advancements In Biocompatible Polymer-Based Nanomaterials For Restorative Dentistry: Exploring Innovations And Clinical Applications : A Literature Review.* *African Journal of Biomedical Research.* 27. 2254-2262.

32. Suda H, Hirabayashi K, Okubo S, et al. The effect of surface treatment on the mechanical stability of dental implants in bone defects. *Clin Oral Implants Res.* 2018; 29(9):881-888.
33. Güncü GN, Beugels J, Meijer HJ, et al. Systemic conditions and their influence on the success of dental implants: A literature review. *Clin Oral Implants Res.* 2018; 29(4):521-530.
34. Friedman M, Zarb GA, Fenton A. A comparison of immediate and delayed loading of dental implants: A retrospective study. *Implant Dent.* 2021; 30(1):15-22.
35. Zhou X, Liao H, Zhang Y, et al. Clinical outcomes of dental implants in patients with osteopenia: A systematic review. *J Prosthet Dent.* 2019; 122(6):505-511.
36. Elian N, Laskin D, Roberts J, et al. Biomechanical properties of dental implants and their effect on osseointegration in compromised bone: A clinical study. *J Oral Maxillofac Surg.* 2019; 77(10):2049-2057.
37. Van Steenberghe D, Gocmen K, Van Landuyt K, et al. Clinical performance of dental implants in patients with osteoporosis: A review. *J Periodontol.* 2019; 90(4):395-405.
38. Iglhaut G, Wichmann M, Buzin K, et al. Osseointegration and loading protocols in compromised bone: Clinical outcomes of implants in patients with osteoporosis. *J Clin Periodontol.* 2020; 47(9):1140-1150.
39. Puppini-Rontani J, Pereira Filho V, Simao R, et al. Systemic conditions and their effects on the healing of dental implants: A clinical review. *J Oral Implantol.* 2019; 45(4):286-292.
40. Aronovich S, Romo J, Misra D, et al. Implant-supported prostheses in patients with high-risk factors: Clinical review of outcomes. *Int J Oral Maxillofac Implants.* 2018; 33(2):289-297.
41. Mavropoulos A, Mouzakis A, Konstantinidis A, et al. The role of local bone density in the success of dental implants: A review of the literature. *J Prosthodont.* 2020; 29(3):185-191.
42. Friberg B, Ekestubbe A, Mellström D, Sennerby L. Brånemark implants and osteoporosis: A clinical exploratory study. *Clin Implant Dent Relat Res* 2001; 3(1):50-6.
43. Silthamptag P, Klineberg I, Austin B, Jones AS. Bone microarchitecture at oral implant sites in ectodermal dysplasia (Ed): A comparison between males and females. *Clin Oral Implants Res* 2012; 23(11):1275-82.
44. Zou D, Wu Y, Wang XD, Huang W, Zhang Z, Zhang Z. A retrospective 3- to 5-year study of the reconstruction of oral function using implant-supported prostheses in patients with hypohidrotic ectodermal dysplasia. *J Oral Implantol* 2014; 40(5):571-80.
45. Merheb J, Temmerman A, Rasmusson L, Kübler A, Thor A, Quirynen M. Influence of skeletal and local bone density on dental implant stability in patients with osteoporosis. *Clin Implant Dent Relat Res* 2016; 18(2):253-60.
46. Wagner F, Schuder K, Hof M, Heuberger S, Seemann R, Dvorak G. Does osteoporosis influence the marginal peri-implant bone level in female patients? A cross-sectional study in a matched collective. *Clin Implant Dent Relat Res.* 2017; 19(4):616-23.
47. Machado M, Wallace C, Austin B, Deshpande S, Lai A, Whittle T, et al. Rehabilitation of ectodermal dysplasia patients presenting with hypodontia: Outcomes of implant rehabilitation part 1. *J Prosthodont Res* 2018; 62(4):473-8.
48. Block MS. Dental implants: The last 100 years. *J Oral Maxillofac Surg.* 2018; 76(1):11-26.
49. Gupta KK, Sawhney H, Nair GK, Gojaniur UC, Kumari A, Kashwani R. A case of benign hard tissue lesion of jaw bone: Report of an enigmatic presentation. *J Adv Sci.* 2024;3(2).
50. Takasaki A, Kawai T, Ishikawa I. Advances in the use of biomaterials for implant surface modifications: A clinical review. *J Clin Med.* 2020; 9(6):1756.

51. Junker R, Dimakis A, Thoneick M, Jansen JA. Effects of implant surface coatings and composition on bone integration: A systematic review. *Clin Oral Implants Res* 2009; 20(suppl 4): 185-206.
52. Pjetursson BE, Tan WK, Lang NP, et al. A systematic review of the survival and complication rates of implants in the posterior maxilla. *Clin Oral Implants Res*. 2019; 30(7):1257-1272.
53. Lemos CA, Verri FR, de Moraes M, et al. Influence of diabetes mellitus on the survival of dental implants: A systematic review and meta-analysis. *J Prosthodont*. 2020; 29(5):363-369.
54. Misch CE, Dietsh F. Implant placement in patients with systemic conditions: A review of literature. *J Periodontol*. 2017; 88(10):991-1000.
55. Tavano L, Bonfiglioli R, et al. Impact of osteoporosis on the survival and success of dental implants: A clinical study. *Clin Oral Implants Res*. 2018; 29(5):1070-1077.
56. Pereira Lima T, Oliveira L, et al. Bone regeneration techniques for compromised bone in dental implantology. *J Clin Periodontol*. 2021; 48(8):1063-1074.
57. Zhang X, Zhang J, Yang C, et al. Osteogenic differentiation of mesenchymal stem cells on titanium implants with different surface modifications. *J Biomed Mater Res A*. 2019; 107(4):892-901.
58. Kao RT, Wang HL. Bone grafting techniques for implant placement in patients with compromised bone: A review of the literature. *Int J Periodontics Restorative Dent*. 2020; 40(6):845-856.
59. Kashwani R, Sawhney H. Dentistry and metaverse: A deep dive into potential of blockchain, NFTs, and crypto in healthcare. *International Dental Journal of Student's Research* 2023;11(3):94-98