

## Platelet-Rich Plasma in Dentistry: Unlocking the Future of Regenerative Healing and Tissue Repair

Dr. Ivianys Darias Otero<sup>1</sup>, Dr. Aditi Pustake<sup>2</sup>, Dr. Puneet Prasher<sup>3</sup>, Dr. Divleen Kaur<sup>4</sup>, Dr. Nikita Duggal<sup>5</sup>, Dr. Dheyaa Al-Alkharsa<sup>6</sup>

1 DMD, Santa Clara, Villa Clara, Cuba. dra.ivianys2015@gmail.com

2 BDS, Maharashtra, India. draditipustake@gmail.com

3 MDS, Punjab, India. docp231189puneet@gmail.com

4 MDS, Punjab, India. divleenkhurana@gmail.com

5 BDS, MHA, Faridkot, Punjab, India. duggalnikita0206@gmail.com

6 BDS, Ternopil, Ukraine. dheyaaakhamees@gmail.com

### KEYWORDS

Platelet-rich plasma (PRP), regenerative dentistry, tissue engineering, wound healing, bone regeneration, implantology, periodontal therapy, oral surgery, temporomandibular joint disorders, endodontics, prosthodontics, orthodontics, soft tissue healing, growth factors, regenerative medicine.

### ABSTRACT

Platelet-rich plasma (PRP) is a promising therapy in regenerative dentistry, known for its ability to accelerate wound healing, enhance bone regeneration, and promote soft tissue repair. Derived from autologous blood, PRP is rich in growth factors and bioactive proteins, contributing to its therapeutic potential. It is widely used in implantology, periodontology, oral surgery, endodontics, and temporomandibular joint (TMJ) disorder management. However, inconsistencies in preparation techniques, platelet concentration, and clinical application challenge its standardization. This review evaluates PRP's biological mechanisms, clinical applications, limitations, and prospects in dentistry. A systematic review of recent studies and clinical trials was conducted, analyzing PRP's impact on osteogenesis, angiogenesis, inflammation modulation, and tissue regeneration. Comparisons with platelet-rich fibrin (PRF) were included to assess relative effectiveness. PRP has shown significant benefits in implant osseointegration, bone graft integration, and soft tissue healing while reducing inflammatory responses and post-operative complications. Key growth factors, including platelet-derived growth factor (PDGF), transforming growth factor-beta (TGF- $\beta$ ), and vascular endothelial growth factor (VEGF), contribute to its regenerative effects. However, variability in preparation protocols and clinical outcomes remains a challenge. While PRP holds great potential in regenerative dentistry, standardized preparation methods, optimized application techniques, and large-scale clinical trials are necessary to ensure consistent results. Future advancements in stem cell integration, biomaterials, and tissue engineering could establish PRP as a cornerstone of next-generation dental therapies.

### Introduction

Platelet-rich plasma (PRP) is an autologous concentrate of platelets obtained from a patient's blood. It has gained prominence in regenerative medicine and dentistry because it stimulates tissue repair, accelerates healing, and promotes cellular regeneration. PRP is particularly valued in oral and maxillofacial surgery, implantology, and periodontal therapy for enhancing wound healing and bone regeneration<sup>1</sup>.

PRP contains a concentrated mix of bioactive proteins and growth factors pivotal in the wound-healing cascade. Due to these biological properties, PRP is now widely used as an adjunct therapy in dental and oral surgeries, enhancing post-surgical recovery and reducing complications<sup>2</sup>.

### **Historical Background**

PRP therapy was first explored in the 1970s for its potential in healing after cardiac surgery. Dr. Knight introduced platelet concentration techniques, which were later expanded into orthopedic and dermatological applications<sup>3</sup>. By the 1990s, its role in regenerative medicine became more pronounced, particularly in treating sports injuries and chronic wound healing<sup>4</sup>.

Dentistry began incorporating PRP in the early 2000s, primarily for bone grafting, implant stabilization, and periodontal tissue healing. As research progressed, its applications broadened to include endodontics, prosthodontics, and maxillofacial reconstruction, making it a vital tool in dental regenerative therapies<sup>5,6</sup>.

### **Mechanism of Action of PRP**

PRP enhances healing by activating a cascade of biological processes that accelerate tissue repair, cell migration, and matrix formation. Once injected or applied to a surgical site, PRP initiates the following mechanisms:

#### Growth Factor Release

Upon activation, platelets undergo degranulation, releasing bioactive molecules that stimulate cellular proliferation and differentiation. Key growth factors include:

- Platelet-derived Growth Factor (PDGF) – Stimulates fibroblast and osteoblast proliferation, enhances angiogenesis, and aids collagen deposition<sup>5</sup>.
- Transforming Growth Factor-Beta (TGF- $\beta$ ) – Regulates immune response, modulates fibrosis, and accelerates wound closure<sup>4</sup>.
- Vascular Endothelial Growth Factor (VEGF) – Induces capillary formation, ensuring vascular support to regenerating tissues<sup>7</sup>.
- Epidermal Growth Factor (EGF) – Promotes epithelial regeneration, reducing the risk of wound dehiscence<sup>6</sup>.
- Insulin-like Growth Factor (IGF) – Enhances bone remodeling and mineralization, essential for implant integration<sup>4</sup>.

#### Cellular Recruitment and Differentiation

PRP attracts and activates several cell types which are fundamental to wound healing:

- Mesenchymal Stem Cells (MSCs) – Differentiate into osteoblasts, fibroblasts, and chondrocytes, depending on the tissue type<sup>4</sup>.
- Fibroblasts – Produce collagen and extracellular matrix proteins essential for soft tissue regeneration<sup>7</sup>.
- Osteoblasts – Facilitate bone matrix formation and mineralization, improving graft stability and osseointegration<sup>2</sup>.

#### Anti-Inflammatory and Antimicrobial Properties

PRP modulates pro- and anti-inflammatory cytokines, reducing excessive inflammation and promoting tissue regeneration. It downregulates TNF- $\alpha$  and IL-1 $\beta$ , while enhancing IL-10, leading to faster resolution of inflammation and minimal scar tissue formation<sup>4</sup>.

Additionally, PRP exhibits antibacterial properties due to the presence of defensins and cathelicidins, which reduce the risk of bacterial contamination in surgical wounds<sup>3</sup>.

#### Angiogenesis and Tissue Vascularization

An essential component of PRP's regenerative ability is angiogenesis, which ensures an adequate blood supply to healing tissues. VEGF, PDGF, and fibroblast growth factor (FGF) promote the formation of microcapillaries, leading to:

- Improved oxygen diffusion into the wound site.
- Efficient metabolic waste removal, reducing oxidative stress.
- Stronger graft vascularization, which enhances bone and soft tissue survival<sup>7,2</sup>.

#### Fibrin Matrix Formation

PRP undergoes fibrin polymerization upon activation, creating a scaffold that supports:

- Cell migration and proliferation, accelerating tissue repair.
- Stabilization of growth factors, ensuring sustained regenerative effects.
- Mechanical support for bone graft materials and implants<sup>3</sup>.

This three-dimensional fibrin network is beneficial in periodontal defect repair and implant site preservation, enhancing bone regeneration and wound stabilization.

#### Enhancement of Soft and Hard Tissue Healing

PRP accelerates both soft (gingival, mucosal) and complex (bone) tissue repair by:

- Reducing healing time post-extraction, implant placement, or grafting<sup>6</sup>.
- Increasing collagen deposition, leading to better tissue strength and elasticity<sup>4</sup>.
- Improving bone density, facilitating stronger implant-bone interactions<sup>2</sup>.
- Research has demonstrated significant improvements in implant survival rates and faster postoperative recovery when PRP is incorporated into surgical protocols<sup>5,6</sup>.

### **Applications of PRP in Dentistry**

PRP in Oral and Maxillofacial Surgery and TMJ Disorders.

Platelet-rich plasma (PRP) is increasingly recognized for its regenerative effects in the fields of oral and maxillofacial surgery, particularly in treating temporomandibular joint (TMJ) disorders. PRP, which is derived from the patient's blood and enriched with growth factors, plays a key role in tissue healing, bone regeneration, and reducing inflammation. This review examines the diverse applications of PRP in oral and maxillofacial surgeries, focusing specifically on its effectiveness in bone healing, soft tissue regeneration, and TMJ disorder management. The findings discussed here are supported by recent studies and clinical research that demonstrate the growing clinical adoption of PRP<sup>1,5,8</sup>.

PRP is a patient-derived concentrate rich in platelets and bioactive proteins. Known for its regenerative capabilities, PRP is used in various oral surgeries such as implantology, bone grafting, and the management of TMJ disorders. Studies show that PRP significantly accelerates the healing process in hard and soft tissues, promoting faster recovery and better clinical outcomes in surgeries like sinus augmentation, alveolar ridge reconstruction, and dental implant integration<sup>5</sup>. Moreover, PRP has positively impacted TMJ disorder management, reducing pain, improving joint function, and supporting cartilage repair through anti-inflammatory and regenerative actions<sup>8</sup>.

### **PRP in Oral and Maxillofacial Surgery**

#### **1. Bone Regeneration and Implantology**

PRP's most prominent use in oral surgery is bone regeneration, frequently combined with bone grafting materials to enhance osteogenesis. The growth factors in PRP stimulate osteoblast activity, which promotes faster bone formation and increases bone density. Several studies have found that when applied along with bone grafts, PRP enhances healing and stability in dental

implants, leading to improved long-term results<sup>1</sup>. Clinical trials have also confirmed that PRP enhances the integration of dental implants and contributes to better bone quality, especially in complex or compromised cases<sup>5</sup>.

## 2. Soft Tissue Healing

In addition to its effects on bone regeneration, PRP plays a vital role in soft tissue healing. Research has proven that PRP accelerates epithelialization, stimulates fibroblast proliferation, and boosts collagen synthesis, all essential for wound healing and reducing postoperative inflammation. A significant reduction in recovery time has been observed in patients undergoing periodontal and maxillofacial procedures with PRP, and its application has been linked to enhanced tissue repair and reduced swelling<sup>5</sup>. PRP also helps control inflammatory responses, leading to decreased postoperative pain and improved overall recovery<sup>1</sup>.

## PRP in Temporomandibular Joint (TMJ) Disorders

### 1. Pain Relief and Joint Function Improvement

TMJ disorders are often associated with chronic pain, inflammation, and restricted jaw movement. Conventional treatments, such as corticosteroid injections and physical therapy provide temporary relief; however, PRP offers a more sustainable solution with longer-lasting benefits. Clinical evidence has shown that PRP injections into the TMJ significantly reduce pain and enhance jaw mobility. For instance, a study by Quijada Gutiérrez et al. (2022) found that patients treated with PRP injections experienced considerable pain relief and improved joint function compared to those receiving traditional therapies<sup>9</sup>. This indicates that PRP is a reliable long-term solution for TMJ disorders.

### 2. Cartilage Regeneration

Cartilage degeneration is a major challenge in treating TMJ disorders, but PRP has shown significant potential in promoting cartilage regeneration. Growth factors such as TGF- $\beta$  and PDGF in PRP stimulate both chondrocyte proliferation and extracellular matrix formation, essential for cartilage repair. Studies have demonstrated that PRP therapy provides symptomatic relief and supports long-term joint health by improving chondrocyte viability and reducing synovial inflammation. PRP is considered an effective treatment option for TMJ osteoarthritis and other degenerative joint diseases<sup>10,11</sup>.

## Challenges and Future Directions

Despite the promising results, several challenges must be addressed before PRP becomes a routine part of clinical practice. One major challenge is the lack of standardization in PRP preparation methods, which can lead to variability in treatment outcomes. Additionally, individual patient responses to PRP therapy, making it crucial to develop more personalized treatment protocols based on specific patient factors<sup>5</sup>. Further research is necessary to optimize PRP preparation techniques, refine its clinical application, and conduct larger, long-term clinical trials. As regenerative medicine continues to evolve, PRP is expected to play an increasingly important role in both surgical and non-surgical treatments for oral and maxillofacial conditions<sup>1</sup>. PRP has emerged as a valuable tool in oral and maxillofacial surgery, particularly in bone regeneration, soft tissue healing, and TMJ disorder management. Its ability to accelerate tissue healing, reduce inflammation, and stimulate cartilage regeneration makes it a valuable adjunct in modern clinical practice. However, additional research is needed to standardize its use and assess its long-term benefits. As evidence continues to accumulate, PRP is likely to play a central role in regenerative therapies for oral and maxillofacial conditions<sup>1,5,8</sup>.

## **PRP in Periodontology and Soft Tissue Healing**

PRP is used for periodontal defect treatment, root coverage procedures, ridge augmentation grafting, guided bone regeneration, sinus lift grafting, and implant surgery. Its clinical application also extends to mandibular and maxillary reconstruction, blepharoplasty, dermal fat grafts, and orthopedic surgery.

Process of PRP in tissue healing and regeneration

In the initial stages of the healing process, platelet-derived growth factors in platelet-enriched plasma attract undifferentiated mesenchymal cells to the fibrin matrix and activate cell division. In this way, they promote tissue regeneration by stimulating the proliferation of connective tissue progenitor cells, fibroblast and osteoblast activity, and angiogenesis. PRP delivers cytokines and growth factors to the site, facilitating rapid regeneration that would not occur with fibrin glue. Because PRP contains platelets, it is free from concerns about transmissible diseases and offers convenience for patients. However, its limitations include a lack of standardization in PRP preparation protocols, variations in storage times for different platelet concentrations, and potential life-threatening coagulopathies, as bovine thrombin can trigger antibodies to clotting factors. (Bovine thrombin can trigger antibodies to clotting factors).

Regenerative periodontal procedures induce regeneration of the alveolar bone and cementum and develop a new functional periodontal ligament. They can be used alone or as an adjunctive material for treating intrabony periodontal defects. PRP or PRF can reduce pocket depth and improve clinical attachment level. However, no beneficial effect was found when they were used along with guided tissue regeneration (GTR).

In a recent systematic review, Plachokova et al.<sup>12</sup> found evidence for the beneficial effects of PRP in the treatment of periodontal defects. Evidence for the beneficial effects of PRP in sinus elevation appeared weak, and no conclusions could be drawn about its other applications in dentistry. Meanwhile, Kotsovilis et al.<sup>13</sup> obtained diverse outcomes for the efficacy of PRP with various therapeutic bioactive agents/procedures, suggesting that the specific selection of agents/procedures combined with PRP is essential. A meta-analysis by Bae et al.<sup>14</sup> indicated sufficient evidence to support using PRP for bone formation in a sinus bone graft. In contrast, its effect on the implant survival was less significant.

### **The role of PRF in periodontal regeneration**

Platelet-rich fibrin is a second-generation platelet concentrate that can enhance both soft and hard tissue healing. Its advantages over platelet-rich plasma include ease of preparation and application, minimal expense, and the absence of biochemical modification. The structural configuration of PRF with cytokine incorporation in fibrin meshes differs from that of PRP. The natural polymerization in PRF results in increased incorporation of circulating cytokines into the fibrin meshes (intrinsic cytokines).

According to Choukron et al., PRF was initially used in implant surgery to enhance the healing properties of the bone<sup>15</sup>. PRF can promote the healing of osseous defects by the following mechanisms, according to Chang et al., PRF encourages the expression of phosphorylated extracellular signal-regulated protein kinase and stimulates the production of osteoprotegerin (OPG), which, in turn, promotes osteoblasts proliferation<sup>16</sup>. Another study by Huang et al. reported that PRF stimulates the osteogenic differentiation of the human dental pulp cells by upregulating osteoprotegerin and alkaline phosphatase expression. PRF also releases growth

factors, such as platelet-derived and transforming growth factors, which promote periodontal regeneration<sup>17</sup>. Chang et al., in a study, reported that PRF stimulates cell proliferation in a specific manner. PRF induces cell proliferation of osteoblasts, periodontal ligament cells, and growth factors during a three-day culture period and suppresses oral epithelial cell growth. These cell type-specific actions may be beneficial for periodontal regeneration. Diss et al.,<sup>18</sup> in a one-year prospective study on osteotome sinus floor elevation using Choukroun's platelet-rich fibrin grafting material, clearly demonstrated that the fibrin matrix of PRF directly promotes angiogenesis. When used as a membrane for guided tissue regeneration or as a grafting material, PRF creates an improved spacemaking effect, facilitating favorable cell events for periodontal regeneration and leading to mineralized tissue formation.

#### Comparison between PRP and PRF

A comparative evaluation between platelet-rich fibrin and platelet-rich plasma for treating three-wall intrabony defects showed a greater bone fill in the PRF-treated group than in the PRP-treated group. The effect of platelet-rich fibrin on human periodontal ligament fibroblasts and application in periodontal infrabony defects was studied by Chang et al., who reported that PRF was found to increase extracellular signal-regulated protein kinase phosphorylation and osteoprotegerin in periodontal ligament fibroblasts and upregulation of alkaline phosphatase activity. PRF, a new generation of platelet concentrate, is a novel step in regenerative periodontal treatment with simplified processing and without biochemical modification. Clinical studies in cooperating PRF for various treatments are encouraging; however, further studies are necessary to support its everyday use in routine dental practice with high clinical efficacy and long-term stability.

#### **PRP in Endodontics and Pediatric Dentistry**

Pulp vitality refers to the health and functionality of the dental pulp, which contains nerves and blood vessels within a tooth. It is essential for maintaining the tooth's sensory function, providing nutrients, and responding to stimuli such as temperature or pressure. Preserving pulp vitality is crucial for the tooth's long-term health and survival, as any damage or infection to the pulp can lead to pain, disease, or even tooth loss if not properly treated.

Pulp necrosis occurs when the pulp dies, which includes the breakdown of blood vessels, lymphatic vessels, and nerve fibers. Conventional endodontic treatments consist of intracanal medications and root-filling materials that are long established and proven effective. However, it is important to recognize that teeth without vital pulp lose their natural defense mechanisms. As a result, they are more prone to fractures and reinfections when treated with conventional treatment protocol<sup>19</sup>. Also, trauma, dental caries, or developmental anomalies can compromise the developing dentition, potentially resulting in pulp necrosis. This condition halts root development, leading to an open apex and thin dentinal walls, which may pose challenges and complications during and after root canal therapy. Apexification has been a conventional approach for preserving non-vital, immature permanent teeth for many years. This procedure involves the formation of a calcified barrier at the open apex. However, it does not promote further root elongation or an increase in root width<sup>20</sup>.

Platelet-rich plasma in regenerative endodontics is a biological scaffold derived from the patient's blood, rich in growth factors, promoting tissue healing and regeneration. It plays a crucial role in pulp-dentin complex regeneration and is particularly useful in procedures like revascularization of necrotic immature teeth and pulp regeneration. PRP is prepared in a gel form by combining PRP -

obtained through the centrifugation of autologous whole blood-with thrombin and calcium chloride. This PRP gel contains a high platelet concentration and a natural fibrinogen level<sup>1</sup>.

A double-blinded parallel randomized controlled trial was implemented to evaluate the potential of PRP and PRF. For twelve months, twenty-five patients were kept on follow-up. PRP demonstrated a slight enhancement in radiographic root length and width, increased periapical bone density, and reduced apical diameter. It was concluded that for necrosed immature teeth, revascularization with PRP served as a suitable alternative to PRF and has demonstrated an excellent prognosis over 12 months<sup>20</sup>.

A triple randomized control trial was done, and the effects of PRP, PRF, and induced bleeding technique on necrotic teeth with open apex. After 12 months, all three groups showed no pain, no signs of reinfection, and no radiographic progression of the pre-existing apical pathosis. Among them, PRP demonstrated superior effectiveness in promoting periapical wound healing compared to PRF and the induced bleeding technique in regenerative endodontic procedures<sup>21</sup>.

Nowadays, a diverse range of materials are available for pulp tissue regeneration. These materials still have some limitations in their use and necessitate additional research and refinement<sup>22</sup>. The limited certainty of the existing evidence highlights the need for further clinical studies to establish the clinical significance of regenerative endodontic procedures as a shift from traditional endodontic practices<sup>19</sup>.

### **Role of platelet-rich plasma in Prosthodontics, Restorative Dentistry and Orthodontics**

Platelet-rich plasma plays a crucial role in both pre-prosthetic and post-prosthetic care, significantly impacting tissue healing, prosthetic function, and overall treatment success. Prosthodontists can significantly improve denture fit, function, and patient comfort by combining PRP with surgical and prosthetic techniques.

PRP is used in vestibuloplasty procedures to deepen the oral vestibule, allowing better adaptation and seating of dentures. It promotes faster healing of soft tissues after vestibuloplasty, reducing post-surgical complications. The use of horizontal platelet-rich fibrin in vestibuloplasty is effective in increasing the vestibule depth and keratinized tissue width. H-PRF in vestibuloplasty has allowed predictable surgery without significant complications. Therefore, it represents an alternative to the traditional FGG that should be seriously considered<sup>23</sup>.

PRP is used in ridge augmentation procedures to increase the vertical height of the alveolar ridge. It accelerates bone regeneration when combined with bone grafts (autografts, allografts, or xenografts), making the ridge more suitable for conventional and implant-supported dentures. In severely resorbed ridges, PRP helps maintain the bone volume, ensuring better support for dentures. The use of PRP in augmenting the severely atrophic posterior maxilla has clear clinical benefits in reducing the healing period of bone maturation, better graft handling, and accelerated soft tissue healing. PRP used with bone graft for sinus floor augmentation to enhance sinus floor while placing implant has found to be effective as there was minimal clinical evidence of crestal bone loss around the implants both clinically and radiographically after 6 months postoperatively<sup>24</sup>.

PRP enhances healing after maxillectomy, improving obturator retention and mucosal adaptation. It can be used with soft tissue grafts to strengthen tissue bed preparation for maxillofacial prostheses. PRP enhances healing and reduces scar formation in cleft lip and palate surgeries. It can be used in secondary alveolar bone grafting to improve bone integration at the cleft site. PRP aids soft tissue healing for patients undergoing radiotherapy after oral cancer treatment and reduces post-radiation oral mucositis and tissue necrosis, enhancing prosthetic outcomes. It improves

osseointegration of zygomatic and extraoral implants (ear, nose, or orbital prosthesis). It is used in titanium mesh reconstruction to enhance healing and reduce complications.

### **Clinical applications in prosthodontics**

PRP + vestibuloplasty → improves denture stability by deepening the vestibule.

PRP + bone grafting → enhances bone height for better denture support and retention.

PRP in implant prosthetics → enhances osseointegration and reduces healing time.

### **PRP in Orthodontics**

Platelet-rich plasma (PRP) is gaining attention in orthodontics due to its regenerative potential and ability to accelerate tissue remodeling. PRP contains growth factors (PDGF, TGF- $\beta$ , VEGF) that stimulate bone remodeling by enhancing osteoclastic and osteoblastic activity. It reduces the overall treatment duration by accelerating bone resorption and deposition. PRP injections near the periodontal ligament can enhance tooth movement in cases requiring rapid space closure. Growth factors in PRP promote cementum repair, potentially preventing further resorption.

It is used in guided tissue regeneration (GTR) and ridge augmentation before or during orthodontic treatment. It can be combined with bone grafts in cases of severe bone loss or open bite corrections. A split-mouth study by Ahmed et al.,<sup>25</sup> concluded that PRP injections when infused with a CaCl<sub>2</sub> activating solution, showed a positive potential to accelerate the rate of tooth movement when injected in the first 2 months. The use of submucosal injection of PRP on orthodontic tooth movement is a clinically feasible and effective technique to accelerate orthodontic tooth movement and at the same time, while preserving the alveolar bone on the pressure side of orthodontic tooth movement, and the optimal dose of PRP for the best clinical performance is 11.0–12.5 fold<sup>26</sup>.

### **PRP in Restorative Dentistry**

PRP is increasingly used in restorative dentistry due to its regenerative properties, promoting healing and enhancing the success of restorative procedures. PRP promotes dentin-pulp complex regeneration in cases of deep caries, traumatic injuries, or pulp exposure. It enhances odontoblast differentiation, leading to the formation of reparative dentin. It can also be used in procedures like direct pulp capping, pulpotomy, and apexogenesis to preserve pulp vitality.

In a controlled animal study, Murat Medan et al.<sup>27</sup> found that PRP can induce or promote inflammatory processes and survive in the wounded rat dental pulp. The use of platelet-rich plasma for regeneration in non-vital, immature permanent teeth suggests that PRP can serve as a successful scaffold for regenerative endodontic treatment<sup>28</sup>.

## **CHALLENGES AND LIMITATIONS OF PRP IN DENTISTRY**

PRP is not universally effective for all dental procedures. It tends to be most successful in areas like periodontal regeneration and bone grafting, but may have limited benefits in soft tissue treatments or for patients with specific health conditions<sup>1</sup>.

### **Variable Outcomes**

PRP's effectiveness can vary from patient to patient, and results are not always predictable. Factors such as the preparation method, the patient's overall health, and the specific dental procedure can influence the success of the treatment.

### **Lack of Standard Protocols**

Currently, there is no universally accepted method for preparing and applying PRP. This lack of standardization can lead to variations in platelet concentration and the overall efficacy of the treatment<sup>29</sup>.

### **Inconsistent Equipment and Techniques**

Different PRP preparation systems may yield inconsistent platelet concentrations, impacting the treatment's regenerative properties and overall quality<sup>29</sup>.

### **High Costs**

The specialized equipment and procedures required to prepare PRP make the treatment expensive, potentially limiting its accessibility for some patients.

### **Need for Specialized Training**

Dentists require specific training to effectively prepare and administer PRP, which increases the cost and complexity of its use.

### **Insufficient Long-Term Data**

While PRP has shown promising results for short-term healing, there is a lack of comprehensive long-term clinical data to validate its effectiveness for dental procedures.

### **Research Gaps**

Further randomized controlled trials and research are necessary to confirm the benefits of PRP and establish standardized protocols for its use in dentistry.

### **Patient-Related Factors**

Certain medical conditions, such as anemia, thrombocytopenia or other blood disorders, can impair platelet function, which may hinder the success of PRP therapy<sup>30</sup>.

### **Risk of Infection**

Although PRP is derived from the patient's blood, the preparation process still carries a small risk of contamination, which could result in infection<sup>29</sup>.

### **Post-Treatment Complications**

Improper PRP administration or technique may lead to complications such as inflammation or insufficient tissue regeneration.

### **Treatment Complexity**

Preparing and applying PRP is a time-consuming process involving blood collection, platelet concentration, and application to the treatment site.

### **Adjunct Treatment**

In many cases, PRP is used as a complementary treatment to other procedures, meaning it may not be sufficient for specific conditions.

## **FUTURE PERSPECTIVES OF PRP IN DENTISTRY**

With ongoing research, standardized PRP preparation and application protocols could be established, resulting in more consistent and reliable results. These universal guidelines could make PRP treatments more accessible and efficient for dental professionals worldwide.

### **Enhanced Efficacy and Targeted Therapies**

PRP may be combined with other biological materials, such as platelet-rich fibrin (PRF), stem cells, or bone morphogenetic proteins, to further enhance its regenerative effects, particularly in complex dental procedures such as bone regeneration and periodontal therapy.

### **Wider Clinical Applications**

Although PRP is currently beneficial for periodontal regeneration, future advancements could expand its use to soft tissue healing, including gingival grafting, mucosal repairs, and wound healing in oral surgeries. Additionally, PRP may be explored in oncology to promote faster healing of soft tissues following oral cancer surgeries or radiotherapy<sup>31</sup>.

### **Technological Advancements**

Developing new automated systems could streamline the PRP preparation process, reducing human error and making it easier for general dental practices to incorporate. Combining PRP with stem cells could significantly enhance regenerative capabilities, improving tissue regeneration and the healing of dental tissues and implants<sup>32</sup>.

### **Regenerative Procedures**

Integrating stem cells, PRP, and other regenerative treatments could lead to groundbreaking solutions for tooth regeneration, bone grafting, and even the potential restoration of lost or damaged teeth.

### **More Comprehensive Research**

Ongoing clinical trials and studies will provide the evidence needed to solidify PRP's long-term benefits and potential in dentistry. This research will guide evidence-based decision-making, helping refine techniques for specific conditions. In the future, PRP, in combination with other regenerative strategies, could help grow new teeth or repair damaged ones. As regenerative dentistry advances, PRP will likely play a central role in regenerating bone and soft tissues, improving outcomes for patients undergoing various dental procedures.

In summary, the future of PRP in dentistry holds great promise. Technology, research, and clinical application advancements are expected to enhance its effectiveness and accessibility. As new evidence emerges and innovative techniques are developed, PRP could become an integral part of more dental treatments, offering better healing, faster recovery times, and potentially revolutionizing procedures like tooth regeneration.

### **Conclusion**

Integrating platelet-rich plasma (PRP) into modern dentistry is significant in regenerative dentistry, offering a biologically driven, minimally invasive approach to enhancing tissue repair, accelerating healing, and improving clinical outcomes. Its versatility spans various specialties, including implantology, oral and maxillofacial surgery, periodontology, endodontics, pediatric dentistry, prosthodontics, and even orthodontics—each benefiting from PRP's ability to modulate inflammation, stimulate cell proliferation, and support both hard and soft tissue regeneration.

Despite its wide-ranging advantages, the clinical integration of PRP is not without its challenges. The lack of standardized preparation protocols, variability in platelet concentrations, and the need for specialized training present obstacles to consistent, reproducible patient outcomes. Moreover, while numerous studies affirm PRP's regenerative potential, long-term, large-scale clinical trials are still required to solidify its efficacy across all its applications.

PRP's future in dentistry appears promising, particularly with the advancement of combination therapies, such as PRP with stem cells, bone morphogenetic proteins (BMPs), or bioengineered scaffolds. These developments could unlock new frontiers in personalized dental medicine, offering more predictable, long-lasting regenerative results. As research continues to refine its application, PRP is poised to transition from an adjunctive treatment to a cornerstone of regenerative dentistry, shaping the next era of patient-centered oral healthcare.

## References:

1. Albanese A, Licata ME, Polizzi B, Campisi G. Platelet-rich plasma (PRP) in dental and oral surgery: from wound healing to bone regeneration. *Immun Ageing*. 2013;10:23.
2. Marrelli M, Scully C, Bianchi A, Mozzati M, Piana G, Gatto R, et al. PRP in dental and oral surgery: wound healing and bone regeneration. *J Oral Maxillofac Surg*. 2012;70(5):1002-1010.
3. Bielecki T, Gazdzik TS, Arendt J, Król W, Wielgus M. Antibacterial effect of platelet gel enriched with growth factors: an in vitro study. *J Bone Joint Surg Br*. 2007;89(3):417-420.
4. Ghanaati S, Booms P, Orłowska A, et al. Advanced platelet-rich fibrin: a new concept for cell-based tissue engineering. *J Oral Implantol*. 2014;40(6):679-689.
5. Marx RE. Platelet-rich plasma: evidence to support its use. *J Oral Maxillofac Surg*. 2004;62(4):489-496. <https://doi.org/10.1016/j.joms.2003.12.015>
6. Del Fabbro M, Bortolin M, Taschieri S, Weinstein R. Is platelet concentrate advantageous for periodontal diseases? *J Periodontol*. 2011;82(8):1100-1111.
7. Wrotniak M, Bielecki T, Gaździk TS. Current opinion about using platelet-rich gel in orthopedics and trauma surgery. *Ortop Traumatol Rehabil*. 2007;9(3):227-238.
8. Anitua E, Fernández-de-Retana S, Alkhraisat MH. Platelet-rich plasma in oral and maxillofacial surgery from the perspective of composition. *Platelets*. 2020;32(2):174-182. <https://doi.org/10.1080/09537104.2020.1856361>
9. Quijada Gutiérrez I, Sábado-Bundó H, Gay-Escoda C. Intra-articular injections of platelet-rich plasma and growth factor-rich plasma with arthrocentesis or arthroscopy in the treatment of temporomandibular joint disorders: A systematic review. *J Stomatol Oral Maxillofac Surg*. 2022;123(5):e327-e335. Available at: <https://pubmed.ncbi.nlm.nih.gov/37341166/>
10. Rajput A, Bansal V, Dubey P, et al. A comparative analysis of intra-articular injection of platelet-rich plasma and arthrocentesis in temporomandibular joint disorders. *J Maxillofac Oral Surg*. 2022;21:168-175. <https://doi.org/10.1007/s12663-020-01351-w>
11. Aishwarya K, Rai M. Injectable platelet-rich fibrin: A promising first-line therapy for temporomandibular joint hypermobility. *J Maxillofac Oral Surg*. 2024;23(4):567-573. <https://doi.org/10.1007/s12663-024-02389-w>
12. Plachokova AS, Nikolidakis D, Mulder J, Jansen JA, Creugers NH. Effect of platelet-rich plasma on bone regeneration in dentistry: a systematic review. *Clin Oral Implants Res*. 2008. [DOI].
13. Kotsovilis S, Markou N, Pepelassi E, Nikolidakis D. The adjunctive use of platelet-rich plasma in the therapy of periodontal intraosseous defects: A systematic review. *J Periodontol Res*. 2010. [DOI].
14. Bae JH, Kim YK, Myung SK. Effects of platelet-rich plasma on sinus bone graft: meta-analysis. *Periodontol*. 2011. [DOI].
15. Dohan DM, Choukroun J, Diss A, et al. Platelet-rich fibrin (PRF): A second-generation platelet concentrate, part I: Technological concept and evolution.
16. Tsai CH, Shen SY, Zhao JH, Chang YC. Platelet-rich fibrin modulates cell proliferation of human periodontally related cells in vitro.
17. Huang FM, Yang SF, Zhao JH, Chan YC. Platelet-rich fibrin increases proliferation and differentiation of human dental pulp cells.

18. Choukroun J, Diss A, Simonpieri A, et al. Platelet-rich fibrin (PRF): A second-generation platelet concentrate, part IV: Clinical effects on tissue healing.
19. Scelza P, Gonçalves F, Caldas I, et al. Prognosis of regenerative endodontic procedures in mature teeth: A systematic review and meta-analysis of clinical and radiographic parameters. *Materials (Basel)*. 2021;14(16):4418. <https://doi.org/10.3390/ma14164418>
20. Rizk HM, Salah Al-Deen MSM, Emam AA. Comparative evaluation of platelet-rich plasma (PRP) versus platelet-rich fibrin (PRF) scaffolds in regenerative endodontic treatment of immature necrotic permanent maxillary central incisors: A double-blinded randomized controlled trial. *Saudi Dent J*. 2020;32(5):224-231. <https://doi.org/10.1016/j.sdentj.2019.09.002>
21. Shivashankar VY, Johns DA, Maroli RK, et al. Comparison of the effect of PRP, PRF, and induced bleeding in the revascularization of teeth with necrotic pulp and open apex: A triple-blind randomized clinical trial. *J Clin Diagn Res*. 2017;11(6):ZC34-ZC39. <https://doi.org/10.7860/JCDR/2017/22352.10056>
22. Kovnatskyi O, Geranin S. Regenerative endodontics: Variety of scaffolds and its benefits for pediatric dentistry. *Ukr Stomatol Zh*. 2023;2(2):107-113. <https://doi.org/10.56569/udj.2.2.2023.107-113>
23. Silvia D, Aldo Z. Horizontal platelet-rich fibrin in vestibuloplasty: A case report. *Cureus*. 2024;16(7):e65862. <https://doi.org/10.7759/cureus.65862>
24. Ziv M, Michael P, Arun K, Jacob L. Platelet-rich plasma for bone graft enhancement in sinus floor augmentation with simultaneous implant placement: Patient series study. *Implant Dent*. 2004;13(1):65-70. <https://doi.org/10.1097/01.ID.0000116454.97671.40>
25. Ahmed E, Fouad El S, Faten E, et al. Effect of platelet-rich plasma on the rate of orthodontic tooth movement: A split-mouth randomized trial. *Angle Orthod*. 2020;90(3):354-361.
26. Liou EJW. The development of submucosal injection of platelet-rich plasma for accelerating orthodontic tooth movement and preserving pressure side alveolar bone. *APOS Trends Orthod*. 2016.
27. Murat M, Ekim O, İhsan F, Burcu S. The inflammatory response of the pulp after direct capping with platelet-rich plasma and enamel matrix derivative: A controlled animal study. *Open J Stomatol*. 2014;4:14-21. <https://doi.org/10.4236/ojst.2014.41004>